

- C. Paraganglioma
- D. Renal cell carcinoma

107. You are seeing a patient with biopsy proven, nonAIDS-related primary CNS lymphoma. What is the best treatment?

- A. Surgical resection followed by XRT and methotrexate chemotherapy
- B. XRT + methotrexate chemotherapy
- C. XRT + temozolomide chemotherapy
- D. Surgical resection followed by XRT and temozolomide chemotherapy

108. What is the approximate 5-year survival of patients with biopsy proven primary CNS lymphoma?

- A. 3 to 4%
- B. 15 to 16%
- C. 30 to 31%
- D. 48 to 49%
- E. 55 to 56%

109. A pituitary tumor is considered a macroadenoma after it has crossed what size threshold?

- A. > 5 mm
- B. > 1 cm
- C. > 2 cm
- D. > 3 cm
- E. > 3.5 cm

110. Approximately what percentage of pituitary adenomas are functioning?

- A. 15%
- B. 35%
- C. 50%
- D. 65%
- E. 80%

111. What type of visual field deficit would a large pituitary macroadenoma cause?

- A. Right homonymous hemianopia
- B. Left superior quadrant hemianopia
- C. Central scotoma
- D. Bitemporal hemianopia

112. What serum marker might help lead you to a diagnosis of suprasellar germinoma?