

D. SDH is more common in infants and elderly patients

26. Which of the following drugs is useful in Ischemic head injury [NMDA antagonist]?

- A. MK – 801
- B. Phencyclidine
- C. Dextromethorphan
- D. Ketamine

27. The most common segment of optic nerve to be injured in closed head injury is

- A. Intracranial
- B. Intraocular
- C. Intracanalicular
- D. Intraorbital

28. Neurofibromatosis type I is associated with

- A. 17 p 11.2
- B. 17 p 12.2
- C. 17 Q 11.2
- D. 17 Q 12.2

29. The best view to demonstrate cervical articular masses is

- A. Pillar's view
- B. Towne's view
- C. Swimmer's view
- D. Caldwell – Luc's view

30. The best view for cervicothoracic junction is

- A. Pillar's view
- B. OMO view [open mouth odontoid view]
- C. Swimmer's view
- D. Towne's view
- E. Caldwell-Luc's view

31. The most common incomplete spinal cord injury is

- A. Posterior cord syndrome
- B. Anterior cord syndrome
- C. Central cord syndrome
- D. Brown-Sequard syndrome

32. The most effective Cervico Thoracic Orthosis for bracing against flexion – extension and rotation is

- A. Yale brace
- B. SOMI brace