

D. Seven liters

20- Which of the following is the LEAST appropriate use of noninvasive positive pressure ventilation (NIPPV)?

- A. Acute respiratory distress syndrome (ARDS)
- B. COPD exacerbation
- C. Obstructive sleep apnea
- D. Multiple sclerosis exacerbation

21- Which of the antibiotics below does NOT augment neuromuscular blockade?

- A. Clindamycin
- B. Neomycin
- C. Streptomycin
- D. Erythromycin

22- Which of the following drugs can prevent tachyarrhythmias in patients with Wolff-Parkinson-White (WPW) syndrome?

- A. Droperidol
- B. Pancuronium
- C. Ketamine
- D. Verapamil

23- Which of the following equals the anti-inflammatory activity of 50 mg of prednisone (Deltasone)?

- A. 100 mg cortisol (Solu-Cortef)
- B. 80 mg methylprednisolone (Solu-Medrol)
- C. 7.5 mg dexamethasone (Decadron)
- D. 4 mg betamethasone (Celestone)

24- Meperidine is contraindicated in patients taking which of the following drugs for Parkinson disease?

- A. Bromocriptine
- B. Trihexyphenidyl (Artane)
- C. Selegiline (Eldepryl)
- D. Amantadine (Symmetrel)

25- Emergence delirium (ED) occurs most often with

- A. Sevoflurane
- B. Desflurane
- C. Ketamine
- D. Propofol

26- The most common reason for patients to rate anesthesia with etomidate as unsatisfactory is

- A. PONV
- B. Pain on injection
- C. Recall of intubation
- D. Postoperative hiccups